

Reaction sequence of blood coagulation

Citation for published version (APA):

Hemker, H. C., & Kahn, M. J. P. (1967). Reaction sequence of blood coagulation. *Nature*, 215, 1201-1202. <https://doi.org/10.1038/2151201a0>

Document status and date:

Published: 01/01/1967

DOI:

[10.1038/2151201a0](https://doi.org/10.1038/2151201a0)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.


Reaction Sequence of Blood Coagulation

IN 1964 Macfarlane¹ proposed a reaction mechanism for the interaction of blood coagulation factors, known as the enzyme cascade. In this hypothesis the blood coagulation factors as they occur in plasma are considered to be zymogens (except factor I; namely, fibrinogen). Coagulation is initiated by the conversion of one of these zymogens into an enzyme as a result of contact with anything other than intact vascular endothelium. This enzyme then acts on another clotting factor to form another active enzyme. All clotting factors thus interact in a given order until prothrombin (factor II) is converted into thrombin. Thrombin then brings about the actual coagulation by converting fibrinogen into fibrin.

The prothrombin converting activity (prothrombinase activity) was first thought to be exerted by activated factor V (ref. 1). As the result of further work it now seems that the prothrombinase activity is displayed by a complex of activated factor X ($f.X_a$), calcium ions (II) and factor V ($f.V$) which are adsorbed together on a phospholipid surface.

The evidence for this concept is of two kinds. (a) It can be shown that $f.X_a$ and $f.V$ are bound by phospholipid micelles, and that calcium ions favour the binding of $f.X_a$. High concentrations of calcium, however, inhibit the binding of $f.V$ (refs. 2-5). (b) The kinetics of the generation of prothrombinase activity in mixtures of $f.X_a$, $f.V$, phospholipid and calcium ions can be explained by assuming that a product of all four reactants is the $f.II$ converting substance. It cannot be explained by a reaction of the cascade type².

This communication shows that evidence can also be provided for the existence of a complex consisting of the clotting factors IX (in activated form) and VIII, and phospholipid, similar to that described in section (a). The interaction between $f.IX_a$ and $f.VIII$ previously has been considered to be of the cascade type, that is



The experiments described in Table 1, however, show a strong similarity between the characteristics of adsorption onto phospholipid of $f.X_a$ and $f.IX_a$, on the one hand, and of $f.V$ and $f.VIII$, on the other. Calcium ions favour the adsorption of $f.X_a$ and $f.IX_a$; excess calcium inhibits the adsorption of $f.V$ and $f.VIII$. Control experiments show that it is the phospholipid that adsorbs the clotting factors. The activity that disappeared from the super-


nantant could be recovered from the resuspended phospholipid sediment. No activity was recorded from the phospholipid in experiments in which no activity was adsorbed from the sample.

Table 1. ADSORPTION OF COAGULATION FACTORS BY PHOSPHOLIPID

Calcium ion conc. (mmolar)	Percentage of clotting factor adsorbed			
	V	VIII	IX	X
1	60	58	27	38
50	<1	41	45	60
100	<1	8	68	76

The figures give the means of three different experiments. Each coagulation factor determination was carried out eight times. The material used as a source of *f.V* and *f.VIII* was barium sulphate adsorbed bovine oxalated plasma. For *f.IX_a* and *f.X_a* oxalated bovine serum was used as a source. To prevent active thrombin being formed from residual amounts of prothrombin, 5 mcg/ml. hirudin was added. The phospholipid used was a suspension of inosithin prepared by homogenizing the crude material in *tris-HCl* buffer 0.02 molar, pH 7.5 containing 0.14 molar NaCl. The reaction mixture consisted of 0.9 ml. plasma or serum, 0.4 ml. CaCl₂ solution in *tris-HCl* 20 mmolar pH 7.5; 0.4 ml. of a suspension of inosithin 5 mg/ml. Half the mixture was centrifuged for 30 min at 100,000*g* at 4° C; in the supernatant the original concentration of phospholipid was restored. The other half of the mixture was stored at 4° C and served as a control. The concentration of the clotting factors in the centrifuged sample was determined and expressed as a percentage of that in the control. The percentage adsorbed was calculated as the amount not recovered in the supernatant. Factors VIII and IX were estimated according to Veltkamp⁹, factors V and X were estimated as described in ref. 10. The concentration of hirudin used did not interfere with the clotting factor determinations.

The role of phospholipid in the interaction of *f.IX_a* and *f.VIII* has so far remained obscure, although it has been proved that this role is a mandatory one⁷. Our experiments strongly suggest that phospholipid interacts with calcium ions, *f.IX_a* and *f.VIII* in a manner similar to the interaction of phospholipid, calcium, *f.X_a* and *f.V*. If this is true, the reaction equation of the interaction would be




Theoretically, it is possible to obtain more evidence for the existence of complex A by investigating the kinetics of the generation of *f.X* converting activity from the purified factors IX_a and VIII, that is, by a procedure analogous to the experiments cited in section (b). Technical difficulties in obtaining the pure factors IX_a and VIII as well as in the measuring of the conversion *f.X* → *f.X_a* seem to prevent this approach in the near future.

The fact that kinetic evidence obtained with impure systems has so far suggested that *f.VIII* is converted enzymatically by *f.IX_a* (ref. 8) remains disturbing. From the results of experiments on the kinetics of the interaction of factors X_a and V, it was evident that if only a limited range of concentrations was investigated, the results obtained simulated that of an enzyme inter-

action. If the experiments were carried over a wide range of concentrations then the stoichiometric interaction is suggested⁶.

We have paid no attention to the form in which *f.V* and *f.VIII* participate in the reactions. By this we do not mean to say that we suppose them to be in the unmodified form in which they occur in intact plasma.

The considerations presented here suggest the following scheme of the intrinsic reaction mechanism of blood coagulation



We thank the Netherlands Foundation for Chemical Research (SON) and the Netherlands Organization for the Advancement of Pure Research (ZWO) for support.

H. C. HEMKER
M. J. P. KAHN

Laboratory of Coagulation Biochemistry,
Haematology Section,
Department of Internal Medicine,
University Hospital,
Leyden.

Received May 10, 1967.

¹ Macfarlane, R. G., *Nature*, **202**, 498 (1964).

² Cole, E. R., Koppel, J. L., and Olwin, J. H., *Thromb. Diathes. Haemorrh.*, suppl. 14, 431 (1965).

³ Esnouf, M. P., and Jobin, F., *Thromb. Diathes. Haemorrh.*, suppl. 17, 103 (1965).

⁴ Jobin, F., and Esnouf, M. P., *Biochem. J.*, **102**, 666 (1967).

⁵ Papahadjopoulos, D., and Hanahan, D. J., *Biochim. Biophys. Acta*, **90**, 436 (1964).

⁶ Hemker, H. C., Esnouf, M. P., Hemker, P. W., Swart, A. C. W., and Macfarlane, R. G., *Nature*, **215**, 248 (1967).

⁷ Schiffman, S., Rapaport, S. I., and Chong, M. M. Y., *Proc. Soc. Exp. Biol. and Med.*, **123**, 736 (1966).

⁸ Biggs, Rosemary, and Macfarlane, R. G., *Thromb. Diathes. Haemorrh.*, suppl. 17, 23 (1965).

⁹ Veltkamp, J., *Thromb. Diathes. Haemorrh.* (in the press, 1967).

¹⁰ Hemker, H. C., and Loelliger, E. A., *Thromb. Diathes. Haemorrh.* (in the press, 1968).